

7

**UNCLASSIFIED**

**AD 4 3 9 3 1 4**

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

64-13

1077-6-28

439314

QUARTERLY REPORT ON NO. 1, 2, 3, 4 - 61-31 Jun 62.  
CONTRACT NO DA 92 + 557 + FEO 2 35675.  
INCLUSIVE DATES October 27, 1961 TO January 31, 1962

~~SUBJECT OF INVESTIGATION~~

EXPLORATION OF NEW CHEMOTHERAPEUTICS  
FOR  
INFECTIOUS DISEASES

~~RESPONSIBLE INVESTIGATOR~~

Dr. Toju Hata, M.D.  
President  
Kitano Institute for Infectious  
Diseases  
Head of Antibiotic Division of  
the said Institute

MAY 14 1964

U.S. Army Research & Development Group (9852) (Far East)  
Office of the Chief of Research and Development  
United States Army  
APO 343

439314

EXPLORATION OF NEW CHEMOTHERAPEUTICS

FOR

INFECTIOUS DISEASES

Fundamental Studies on Protomycin, an Antiamoebic  
Antibiotic and Cephalomycin, an Antiviral Anti-  
biotic

Toju Hata,

Ryozo Sugawara, Akihiro Matsumae

and

Hiroshi Yamamoto

Kitasato Institute  
for  
Infectious Diseases

Tokyo, Japan


The main products may be summerized as follows:

3,5-dimethyl-3,5-heptadiene-2-one (III) and acetone (IV) were proved as volatile ketones arising from alkaline degradation of pretomycin.

Formaldehyde (VII) and acetaldehyde (VIII) were proven by ozonolysis of protomyoin.

The oxime of tetrahydropyromycin, although it remained oil, were treated with conc.  $H_2SO_4$  to induce Beckmann rearrangement and thereafter steam distilled from acidic solution. An acid with molecular formula of  $C_{10}H_{19}COOH$  was obtained as p-phenylazophenacyl ester (XI).

These products from (I) through (XII) were found to be explained well by the chemical structure:



### Tentative Structure of Protomycin

Cephalomycin, an antiviral antibiotic, was separated into two fractions as described in p. 17 of the final report (2). Each of these fractions gave a single spot when tested on electrophoretic paperchromatography.

The fraction A migrated as albumin and Fraction B as gamma globulin.

The activity was higher in the former than in the latter.

The result is to be made publish at the Symposium for the Antiviral substance, March 3, 1962.